

A First Encounter: Fundamentals of Brain Structure and Function

How Your Brain Works - Week 1

Prof. Jan Schnupp

wschnupp@cityu.edu.hk

HowYourBrainWorks.net



Course Objective

You are your brain.

Every sight you ever see, every thought you ever think, every desire you ever feel, every dream you ever dream, all your decisions and movements are just manifestations of brain activity.

They should be understandable and explainable in terms of brain organisation (anatomy) and function (physiology).

In this course we aim to give you a sense of the science that underpins these remarkable claims.

Course Outline

- See the Course Schedule posted on Canvas
- There will be weekly lectures and tutorials.
- Tutorials WILL FREQUENTLY CONTAIN QUIZZES to test your knowledge and understanding of the preceding lectures. The quizzes count toward continuous assessment. So make sure you revise your lecture notes before each tutorial and don't hesitate to ask the relevant TA (listed in course schedule) for advice.

Dos and Don'ts

DO ask questions!



DO ask questions in the breaks during lectures!

DO ask questions in the tutorials!

DO send questions to the TAs by email!

DON'T just sit there if you don't understand something. Ask questions!

A First Encounter



- The brain together with the spinal cord and the cranial nerves form the **central nervous system (CNS)**. It controls numerous conscious and unconscious functions of your body, some as simple as reflexes, others as complex as dreams.

The Oldest Brian Functions: “Reflexes”



- In his 1664 “Treatise of Man”, Rene Descartes correctly hypothesized that nerves are conductors of information from the skin and sense organs to the brain and back from the brain to the muscles.
- **The central nervous system evolved as a device made of nerve cells that processes information to convert sensory input into appropriate output.**

Spinal Chord

- is the oldest part of the central nervous system. It takes in information from sensory neurons at the back (the “dorsal nerve roots”), and sends information back out through nerves at the front (the “ventral roots”).





Fig. 1

A Simple Reflex Arc (Patellar Reflex)




- Reflex arcs provide very simple examples of neural information processing.
- A tap on the patellar tendon stretches the quadriceps muscle.
- This activates sensory stretch receptor neurons (“dorsal root ganglion cells”) which connect to motor neurons in the ventral horn of the spinal cord. The motor neurons cause the quadriceps muscle to contract.
- At the same time inhibitory interneurons in the ventral horn tell motor neurons which innervate the hamstrings to relax.
- Increased load on the muscle thus results in increased muscle tension, (as well as relaxation of the antagonist muscle)

Reflex Arcs

- The patellar reflex is an incredibly simple neural circuit (“monosynaptic”) which helps with postural adjustment (resist a sudden bending of the knee).
- There are many more complex reflexes (e.g. withdrawal reflexes) and neural control structures (e.g. central pattern generators which produce rhythmic locomotive movement) that “are bolted on top” of these simple reflexes.
- Note that in this reflex, sensory information enters the central nervous system at the back, and motor outputs are computed and sent out of the front. This is a general pattern of organization that is also true to an extent in higher parts of the brain.
- Nerve cells which carry information toward higher processing centres are called **afferents**. Nerve cells that carry signals toward motor output structures are called **efferents**.


Loops upon Loops



- It is useful to imagine the brain as having evolved by layering more and more complex input-output (“reflex”) loops on top of existing ones.
- Charles Sherrington (Nobel Prize winning Physiologist of the early 1900s) imagined that all brain function might be explainable in terms of potentially hugely complex, interlinked and hierarchically organized reflex arcs which interact.
- More recently, brain researchers feel that Sherrington’s view underestimates the importance of innate drives as well as long term memories in shaping perception and motivation (“cognitive” neuroscience)

Brain Evolution

Over 100s of millions of years, more and more circuits are added to the brain and expand.



Take-home Message

The central nervous system comprises a hierarchy of circuits which evolved gradually.

Some circuits are more primitive than others, and there may be multiple circuits for any one role.

For example, the older *superior colliculus* (aka *optic tectum*) and the newer *visual cortex* both help to make you see. Ancient spinal chord reflex arcs as well as cerebellar and motor cortical circuits all control the way you move.

These circuits are all active in parallel, somewhat independently of each other. It is surprising that we can ever be “of one mind” about anything.


Neurons

The brain's circuits are made of nerve cells or neurons. A human brain contains in the order of 10 billion of them.

Neurons differ greatly in shape and size, but on average the cell bodies of human neurons are roughly about 20 microns across. So in a volume of just $1 \times 1 \times 1$ mm you could fit about $50 \times 50 \times 50 = 125,000$ neurons.


Let's take a closer look.

A “typical” neuron




- Most neurons, or nerve cells, have long **axons** that serve as their “output cable” through which they transmit information.
- Signals are transmitted along axons using **electrical** signals.
- At the end of axons, **synapses** serve to use **chemical signals** to transmit the information on to another neuron (or to a muscle fibre or gland).
- Many neurons also have **dendrites** which receive chemical signals from many (often several thousand!) other neurons and serve as “input cables”.

Neurons as Signal Processing Devices



Golgi stained
“pyramidal”
neuron



Schematic
drawing

Dendrites

Soma


Axon

The Golgi stain reveals different cellular subdivisions in neurons

- *Dendrites* are the main “input cables”
- The cell body, or “soma”, contains the nucleus. Proteins are synthesized in the soma.
- (“Soma” is just a Greek word for “body”)
- There is typically only one “output cable”, the *axon*.
- Axons can project over relatively long distances (up to several meters)


Cortical Pyramidal Neuron

Cerebellar Purkinje Neuron




- Neurons tend to have several “dendrites” (input cables) but only one “axon” (output cable) each.

Communication Between Neurons



- The axon of one neuron typically contacts the dendrites or somata of many others
- The connections between axons and dendrites are known as ***synapses***.
- Synapses are chemical switches: *neurotransmitters* control the opening of ion channels, and allow electrical currents to flow into or out of the receiving cell.
- Synaptic currents from as many as 10,000 synapses are *integrated* (i.e. combined) to determine the voltage at the axon hillock
- The voltage at the axon hillock determines the rate of “nerve impulses” (*action potentials*) sent down the axon to other neurons.

Neural Networks as Computing Devices




- Neurons often receive highly convergent inputs (many synapses contact their many dendrites) and they send out highly divergent outputs (through many axon branches). Consequently, neurons form densely connected networks.
- The synapses between neurons can be “plastic”, meaning that the connection strength of individual synapses can change. This allows the connection patterns in neural network to change in a way that learns new “associations” between the pieces of information represented by the activity of individual neurons.
- Artificial neural networks (ANNs) are inspired by brain research and led to breakthroughs in artificial intelligence, such as machine translation, “go” world championship, self driving cars etc...

Your brain in numbers

- You have about 10 billion neurons in your brain. (And there are yet more neurons in your spinal cord and even in your gut)
- Most neurons receive and send out thousands of “synaptic connections”.
- There are tens of thousands of times more synapses in a human brain than there are seconds in a human lifetime.

Neurons Are Not the Only Cell Type in the Nervous System



- In fact, there are a lot more *glial* cells than neurons in the nervous system.
- They are **not** thought to carry sensory or motor signals themselves.
- There is a diversity of glial cells:
- **Astrocytes** important for the metabolic regulation of the nervous system,
- **Oligodendrocytes & Schwann** cells provide the myelin sheath surrounding axons,
- **Microglia** are involved in immunological reactions to damage in the nervous system.

“White Matter” and “Gray Matter”

Brain

Spinal Cord

Gray matter

White matter








Fig. 1


- Gray matter contains many nerve cell bodies (neurons)
- White matter contains mostly nerve fibers (axons)

Decussations




- Curiously, many of the input and output pathways to and from the brain are “crossed”, so that the left brain hemisphere predominantly processes information from and for the right side of the body and vice-versa.
- Crossing points of nerve tracts responsible for this are called “decussations”. Prominent examples include the optic chiasm, or the “decussation of the pyramids” in the cortico-spinal tract.

Brain Hemispheres




- Much like other systems of the body (skeleton, lungs, sensory organs) the brain is (approximately) mirror symmetric about the midline (the “mid-sagittal plane”, as anatomists would say).

Beware of Brain Hemisphere Myths




- While there is some hemispheric specialisations (most prominently, the left brain is usually much more involved in language processing), **many popular notions of “left brain vs right brain” faculties are oversimplified to the point of being wrong.**

Two Brain Hemispheres, but only one Self



- Rene Descartes found the brain hemispheres odd. How can I have only one unified conscious awareness in a brain that appears to be split into two hemispheres?
- He therefore suggested that consciousness should reside in the one part of the brain he could find that had no discernible midline symmetry: the pineal gland.
- A clever idea, but he was wrong. We now know that the pineal secretes hormones, including melatonin, which makes you sleepy at night.


Commissures



- There are many nerve fibre bundles providing effective communication between the two hemispheres of the brain.
- Fibre bundles connecting one brain hemisphere to the other are called “**commissures**”. The most prominent commissure is the **corpus callosum** which connects the two hemispheres of the cerebral cortex.

Let's take a short break

Anatomical Directions



Rostral Toward face

Caudal Toward the tail

Ventral Toward the belly

Dorsal Toward the back


Medial Toward midline

Lateral Toward the side


Inferior Lower, below

Superior Higher, above

Parts of the Brain




Parts of the Brain: The Meninges




- The **meninges** are a three layered skin covering the central nervous system. It's layers are the *dura* (the “hard” one), the *arachnoid* (the “spiderous” one) and the *pia* (the “soft” one).
- Meningitis is when the meninges become infected and inflamed. Usually a bacterial disease (meningococcus) which can be very serious.
- The arachnoid also reabsorbs cerebro-spinal fluid which is generated in the brain's ventricles.

Parts of the Brain: Ventricles




- The brain also contains a number of cavities known as “ventricles” which are filled with **cerebro-spinal fluid (CSF)**.
- The CSF is actually secreted by the choroid plexus, and then circulates through the ventricles to the spinal cord and finally to the meninges, where it gets reabsorbed by the arachnoid.
- Blockage of CSF circulation can lead to severe swelling of the ventricles, known as **hydrocephalus** (Greek for “water head”)

Parts of the Brain: The Blood-Brain Barrier





- The capillaries (tiny blood vessels) in most of the body are slightly porous, allowing nutrients to leak from the blood serum to the surrounding tissue.
- Brain capillaries have no pores, and the exchange of substances between brain and blood is actively regulated by carrier proteins produced by astrocytes.
- This ensures that certain substances which are abundant in food and which could interfere with the brain's signalling mechanisms (salts, amino acids) cannot just rush into the brain after a meal. It also keeps many water soluble drugs and poisons out of the brain.
- However, fat soluble substances, including alcohol or cannabis, can diffuse through the blood-brain barrier.

Cortex



- Cortex is subdivided into two hemispheres, each comprising four “lobes”
 - Frontal lobe (movement, cognition)
 - Parietal lobe (touch, spatial orientation and attention)
 - Occipital lobe (vision)
 - Temporal lobe (hearing, object recognition, memory)
- Human cortex has prominent ridges (gyri) and grooves (sulci)


Cortex




- Think of cortex of a “sheet” of nerve cells, about 2 mm thick, covering an area of $2,500 \text{ cm}^2$, “crumpled up” to fit into the skull.
- Each mm^3 of the cortical sheet contains about 100,000 neurons.
- The neurons come in different “types”, so called pyramidal or stellate neurons, or basket or granule cells, for example, and they are arranged in six “layers”, where different layers have somewhat different functions and connectivity.

“Brodmann” Areas

- Corbinian Brodmann, 1909.
- 52 different areas delineated based on “cytoarchitectonic” differences of different regions of cortex (from just a single subject).




“Brodmann” Areas



- Brodmann's cytoarchitectonic areas correlate very well with functional specializations identified later. For example, areas 1-3 are “primary somatosensory cortex”, area 17 is primary visual cortex.
- Area 17 (visual cortex) has a very prominent layer IV, Area 4 (primary motor cortex) has almost no layer 4.

Just as in the spinal cord, where sensory information arrives at the back and motor output goes out the front, in the cortex too sensory processing areas are mostly found in the back (the posterior half), and motor control and decision making structures are more in the front (anterior half).


Parts of the Brain: Thalamus



Thalamus


- The Thalamus is an “obligatory relay” between sensory input and cortex. That means (with the exception of smell), sensory afferents from all the senses synapse in thalamus and thalamic neurons then connect to cortex.
- But cortex also sends nerve fibres back down to thalamus, perhaps to send signals of expectation or selective attention to thalamic relay neurons, so that thalamus sends only the most relevant information to cortex.
- Thalamic activity is also thought to be very important in mediating sleep and in directing attention.

Parts of the Brain: Hypothalamus



- Greek “hypo” means “below”.
- The hypothalamus is an ancient brain structure that controls many important bodily functions, such as appetite, thirst, blood pressure, body temperature, sleep-wake (circadian) rhythms, the regulation of the levels of growth hormones, stress hormones and sex hormones, etc...


Parts of the Brain: Pituitary Gland



Pituitary


- The pituitary is perhaps the most important hormone secreting glands of the body.
- It is connected to the hypothalamus via the pituitary stalk, and the hypothalamus directly controls it's function.
- It gets inputs from the hypothalamus to control hormone levels involved in functions as diverse as reproduction, growth, thirst or stress levels.

Parts of the Brain: Basal Ganglia




- Clumps of (often many thousands of) neurons can be referred to as “ganglia” or “nuclei”.
- The Basal Ganglia form feedback circuits, which start in frontal and prefrontal parts of cortex and pass through the basal ganglia to thalamus and back to cortex.
- Parts of the Basal Ganglia include the Caudate and Putamen (together known as the Striatum), the Globus Pallidus, the Subthalamic Nucleus and the Substantia Nigra.
- The Basal Ganglia are hugely important in action selection, habits, even addiction.

Parts of the Brain: Cerebellum



- The cerebellum (Latin for “little brain”) is another brain area involved in motor control.
- It receives diverse sensory inputs from all over the brain and is thought to learn to associate that with motor output patterns in order to make motor execution “smooth” and automatic after practice.
- Damage to the cerebellum makes movements slow and clumsy (ataxia and apraxia), or can cause disturbed balance.
- Its role in cognition and emotion is hotly debated.

Parts of the Brain: The Limbic System



- A group of structures situated around the “edge” (Latin: “limbus”) of the cortex are commonly called the “limbic system”. Key members of the limbic system include the amygdala, hypothalamus, thalamus, hippocampus and cingulate cortex.
- Limbic structures have long been suspected to be both “old” in evolutionary terms as well as important in “emotional” responses, including fear, anger, pleasure and arousal.

Key Summary “Take Home” Messages

- The nervous system controls your muscles and your glands. It controls a vast amount of stuff, from how moist to keep your eyeballs through your tear glands to deciding which courses you will choose next semester, or who you fall in love with.
- The brain's parts are subdivided in symmetric, hemispheric pairs and are richly interconnected by fiber bundles.
- They contain huge numbers of neurons which process information using partly electrical and partly chemical signalling, which we will study in great detail in the next lecture.
- The brain consists of many parts which are thought to have evolved sequentially, with newer, more “modern” structures both adding more sophisticated levels of control which incorporate better analysis of sensory information and which can learn from experience.
- Example of old structures: spinal cord with its reflexes and simple motor patterns, or hypothalamus with its control of hormones and basic drives.
- Example of newer structure: (neo)cortex which is enormous in humans and which is needed for analytical or creative thinking or for language.
- At any one moment in time, your cortex and your limbic structures may have quite different “opinions” about a particular situation.